

PCTWORLD INTELLECTUAL
PROPERTY ORGANIZATION

INTERNATIONAL APPLICATION PUBLISHED

WO 9602555A1

(51) International Patent Classification 6 :

C07H 21/00, C12Q 1/68, A61K 31/70

A1

(43) International Publication Date: 1 February 1996 (01.02.96)

(21) International Application Number: PCT/US95/01570

(22) International Filing Date: 7 February 1995 (07.02.95)

(30) Priority Data:

08/276,358

15 July 1994 (15.07.94)

US

(71) Applicant: THE UNIVERSITY OF IOWA RESEARCH
FOUNDATION [US/US]; 214 Technology Innovation
Center, Oakdale Research Campus, Iowa City, IA 52319
(US).(72) Inventor: KRIEG, Arthur, M.; 890 Park Place, Iowa City, IA
52319 (US).(74) Agents: ARNOLD, Beth, E. et al.; Lahive & Cockfield, 60
State Street, Boston, MA 02109 (US).(81) Designated States: AU, CA, JP, European patent (AT, BE,
CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT,
SE).

Published

With international search report.

(54) Title: IMMUNOMODULATORY OLIGONUCLEOTIDES

(57) Abstract

Oligonucleotides containing unmethylated CpG dinucleotides and therapeutic utilities based on their ability to stimulate an immune response in a subject are disclosed. Also disclosed are therapies for treating diseases associated with immune system activation that are initiated by unmethylated CpG dinucleotides in a subject comprising administering to the subject oligonucleotides that do not contain unmethylated CpG sequences (i.e. methylated CpG sequences or no CpG sequence) to outcompete unmethylated CpG nucleic acids for binding. Further disclosed are methylated CpG containing dinucleotides for use in antisense therapies or as in vivo hybridization probes, and immunoinhibitory oligonucleotides for use as antiviral therapeutics.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

IMMUNOMODULATORY OLIGONUCLEOTIDES

5

Government Support

The work resulting in this invention was supported in part by National
Institute of Health Grant No. R29-AR42556-01. The U.S. Government may therefore be
10 entitled to certain rights in the invention.

Background of the Invention

DNA binds to cell membrane and is internalized

15 In the 1970's, several investigators reported the binding of high molecular
weight DNA to cell membranes (Lerner, R.A., W. Meinke, and D.A. Goldstein. 1971.
"Membrane-associated DNA in the cytoplasm of diploid human lymphocytes". *Proc. Natl.
Acad. Sci. USA* 68:1212; Agrawal, S.K., R.W. Wagner, P.K. McAllister, and B. Rosenberg.
1975. "Cell-surface-associated nucleic acid in tumorigenic cells made visible with platinum-
20 pyrimidine complexes by electron microscopy". *Proc. Natl. Acad. Sci. USA* 72:928). In
1985 Bennett et al. presented the first evidence that DNA binding to lymphocytes is similar
to a ligand receptor interaction: binding is saturable, competitive, and leads to DNA
endocytosis and degradation (Bennett, R.M., G.T. Gabor, and M.M. Merritt. 1985. "DNA
binding to human leukocytes. Evidence for a receptor-mediated association, internalization,
25 and degradation of DNA". *J. Clin. Invest.* 76:2182). Like DNA, oligodeoxyribonucleotides
(ODNs) are able to enter cells in a saturable, sequence independent, and temperature and
energy dependent fashion (reviewed in Jaroszewski, J.W., and J.S. Cohen. 1991. "Cellular
uptake of antisense oligodeoxynucleotides". *Advanced Drug Delivery Reviews* 6:235;
Akhtar, S., Y. Shoji, and R.L. Juliano. 1992. "Pharmaceutical aspects of the biological
30 stability and membrane transport characteristics of antisense oligonucleotides". In: Gene
Regulation: Biology of Antisense RNA and DNA. R.P. Erickson, and J.G. Izant, eds. Raven
Press, Ltd. New York, pp. 133; and Zhao, Q., T. Waldschmidt, E. Fisher, C.J. Herrera, and
A.M. Krieg., 1994. "Stage specific oligonucleotide uptake in murine bone marrow B cell
precursors". *Blood*, 84:3660). No receptor for DNA or ODN uptake has yet been cloned, and
35 it is not yet clear whether ODN binding and cell uptake occurs through the same or a different
mechanism from that of high molecular weight DNA.

Lymphocyte ODN uptake has been shown to be regulated by cell activation.
Spleen cells stimulated with the B cell mitogen LPS had dramatically enhanced ODN uptake

in the B cell population, while spleen cells treated with the T cell mitogen Con A showed enhanced ODN uptake by T but not B cells (Krieg, A.M., F. Gmelig-Meyling, M.F. Gourley, W.J. Kisch, L.A. Chrisey, and A.D. Steinberg. 1991. "Uptake of oligodeoxyribonucleotides by lymphoid cells is heterogeneous and inducible". *Antisense Research and Development* 1:161).

Immune effects of nucleic acids

Several polynucleotides have been extensively evaluated as biological response modifiers. Perhaps the best example is poly (I,C) which is a potent inducer of IFN production as well as a macrophage activator and inducer of NK activity (Talmadge, J.E., J. Adams, H. Phillips, M. Collins, B. Lenz, M. Schneider, E. Schlick, R. Ruffmann, R.H. Wiltout, and M.A. Chirigos. 1985. "Immunomodulatory effects in mice of polyinosinic-polycytidylic acid complexed with poly-L-lysine and carboxymethylcellulose". *Cancer Res.* 45:1058; Wiltout, R.H., R.R. Salup, T.A. Twilley, and J.E. Talmadge. 1985. "Immunomodulation of natural killer activity by polyribonucleotides". *J. Biol. Resp. Mod.* 4:512; Krown, S.E. 1986. "Interferons and interferon inducers in cancer treatment". *Sem. Oncol.* 13:207; and Ewel, C.H., S.J. Urba, W.C. Kopp, J.W. Smith II, R.G. Steis, J.L. Rossio, D.L. Longo, M.J. Jones, W.G. Alvord, C.M. Pinsky, J.M. Beveridge, K.L. McNitt, and S.P. Creekmore. 1992. "Polyinosinic-polycytidylic acid complexed with poly-L-lysine and carboxymethylcellulose in combination with interleukin 2 in patients with cancer: clinical and immunological effects". *Canc. Res.* 52:3005). It appears that this murine NK activation may be due solely to induction of IFN β secretion (Ishikawa, R., and C.A. Biron. 1993. "IFN induction and associated changes in splenic leukocyte distribution". *J. Immunol.* 150:3713). This activation was specific for the ribose sugar since deoxyribose was ineffective. Its potent in vitro antitumor activity led to several clinical trials using poly (I,C) complexed with poly-L-lysine and carboxymethylcellulose (to reduce degradation by RNase) (Talmadge, J.E., et al., 1985. cited *supra*; Wiltout, R.H., et al., 1985. cited *supra*; Krown, S.E., 1986. cited *supra*; and Ewel, C.H., et al., 1992. cited *supra*). Unfortunately, toxic side effects have thus far prevented poly (I,C) from becoming a useful therapeutic agent.

Guanine ribonucleotides substituted at the C8 position with either a bromine or a thiol group are B cell mitogens and may replace "B cell differentiation factors" (Feldbush, T.L., and Z.K. Ballas. 1985. "Lymphokine-like activity of 8-mercaptoguanosine: induction of T and B cell differentiation". *J. Immunol.* 134:3204; and Goodman, M.G. 1986. "Mechanism of synergy between T cell signals and C8-substituted guanine nucleosides in humoral immunity: B lymphotropic cytokines induce responsiveness to 8-mercaptoguanosine". *J. Immunol.* 136:3335). 8-mercaptoguanosine and 8-bromoguanosine also can substitute for the cytokine requirement for the generation of MHC restricted CTL

(Feldbush, T.L., 1985. cited *supra*), augment murine NK activity (Koo, G.C., M.E. Jewell, C.L. Manyak, N.H. Sigal, and L.S. Wicker. 1988. "Activation of murine natural killer cells and macrophages by 8-bromoguanosine". *J. Immunol.* 140:3249), and synergize with IL-2 in inducing murine LAK generation (Thompson, R.A., and Z.K. Ballas. 1990. "Lymphokine-activated killer (LAK) cells. V. 8-Mercaptoguanosine as an IL-2-sparing agent in LAK generation". *J. Immunol.* 145:3524). The NK and LAK augmenting activities of these C8-substituted guanosines appear to be due to their induction of IFN (Thompson, R.A., et al. 1990. cited *supra*). Recently, a 5' triphosphorylated thymidine produced by a mycobacterium was found to be mitogenic for a subset of human $\gamma\delta$ T cells (Constant, P., F. Davodeau, M.-A. Peyrat, Y. Poquet, G. Puzo, M. Bonneville, and J.-J. Fournie. 1994. "Stimulation of human $\gamma\delta$ T cells by nonpeptidic mycobacterial ligands" *Science* 264:267). This report indicated the possibility that the immune system may have evolved ways to preferentially respond to microbial nucleic acids.

Several observations suggest that certain DNA structures may also have the potential to activate lymphocytes. For example, Bell et al. reported that nucleosomal protein-DNA complexes (but not naked DNA) in spleen cell supernatants caused B cell proliferation and immunoglobulin secretion (Bell, D.A., B. Morrison, and P. VandenBygaart. 1990. "Immunogenic DNA-related factors". *J. Clin. Invest.* 85:1487). In other cases, naked DNA has been reported to have immune effects. For example, Messina et al. have recently reported that 260 to 800 bp fragments of poly (dG) \bullet (dC) and poly (dG \bullet dC) were mitogenic for B cells (Messina, J.P., G.S. Gilkeson, and D.S. Pisetsky. 1993. "The influence of DNA structure on the in vitro stimulation of murine lymphocytes by natural and synthetic polynucleotide antigens". *Cell. Immunol.* 147:148). Tokunaga, et al. have reported that dG \bullet dC induces γ -IFN and NK activity (Tokunaga, S. Yamamoto, and K. Namba. 1988. "A synthetic single-stranded DNA, poly(dG,dC), induces interferon- α/β and - γ , augments natural killer activity, and suppresses tumor growth" *Jpn. J. Cancer Res.* 79:682). Aside from such artificial homopolymer sequences, Pisetsky et al. reported that pure mammalian DNA has no detectable immune effects, but that DNA from certain bacteria induces B cell activation and immunoglobulin secretion (Messina, J.P., G.S. Gilkeson, and D.S. Pisetsky. 1991. "Stimulation of in vitro murine lymphocyte proliferation by bacterial DNA". *J. Immunol.* 147:1759). Assuming that these data did not result from some unusual contaminant, these studies suggested that a particular structure or other characteristic of bacterial DNA renders it capable of triggering B cell activation. Investigations of mycobacterial DNA sequences have demonstrated that ODN which contain certain palindrome sequences can activate NK cells (Yamamoto, S., T. Yamamoto, T. Kataoka, E. Kuramoto, O. Yano, and T. T. kunaga. 1992. "Unique palindromic sequences in synthetic ligonucleotides are required to induce INF and augment INF-mediated natural killer activity". *J. Immunol.* 148:4072; Kuramoto, E., O.

Yano, Y. Kimura, M. Baba, T. Makino, S. Yamamoto, T. Yamamoto, T. Kataoka, and T. Tokunaga. 1992. "Oligonucleotide sequences required for natural killer cell activation". *Jpn. J. Cancer Res.* 83:1128).

- 5 Several phosphorothioate modified ODN have been reported to induce in vitro or in vivo B cell stimulation (Tanaka, T., C.C. Chu, and W.E. Paul. 1992. "An antisense oligonucleotide complementary to a sequence in Iy2b increases γ 2b germline transcripts, stimulates B cell DNA synthesis, and inhibits immunoglobulin secretion". *J. Exp. Med.* 175:597; Branda, R.F., A.L. Moore, L. Mathews, J.J. McCormack, and G. Zon. 1993. "Immune stimulation by an antisense oligomer complementary to the rev gene of HIV-1". *Biochem. Pharmacol.* 45:2037; McIntyre, K.W., K. Lombard-Gillooly, J.R. Perez, C. Kunsch, U.M. Sarmiento, J.D. Larigan, K.T. Landreth, and R. Narayanan. 1993. "A sense phosphorothioate oligonucleotide directed to the initiation codon of transcription factor NF- κ β T65 causes sequence-specific immune stimulation". *Antisense Res. Develop.* 3:309; and
- 10 Pisetsky, D.S., and C.F. Reich. 1993. "Stimulation of murine lymphocyte proliferation by a phosphorothioate oligonucleotide with antisense activity for herpes simplex virus". *Life Sciences* 54:101). These reports do not suggest a common structural motif or sequence element in these ODN that might explain their effects.
- 15

- 20 The CREB/ATF family of transcription factors and their role in replication
- The cAMP response element binding protein (CREB) and activating transcription factor (ATF) or CREB/ATF family of transcription factors is a ubiquitously expressed class of transcription factors of which 11 members have so far been cloned (reviewed in de Groot, R.P., and P. Sassone-Corsi: "Hormonal control of gene expression: Multiplicity and versatility of cyclic adenosine 3',5'-monophosphate-responsive nuclear regulators". *Mol. Endocrin.* 7:145, 1993; Lee, K.A.W., and N. Masson: "Transcriptional regulation by CREB and its relatives". *Biochim. Biophys. Acta* 1174:221, 1993.). They all belong to the basic region/leucine zipper (bZip) class of proteins. All cells appear to express one or more CREB/ATF proteins, but the members expressed and the regulation of mRNA
- 25 splicing appear to be tissue-specific. Differential splicing of activation domains can determine whether a particular CREB/ATF protein will be a transcriptional inhibitor or activator. Many CREB/ATF proteins activate viral transcription, but some splicing variants which lack the activation domain are inhibitory. CREB/ATF proteins can bind DNA as homo- or hetero- dimers through the cAMP response element, the CRE, the consensus form of which is the unmethylated sequence TGACGTC (binding is abolished if the CpG is methylated) (Iguchi-Ariga, S.M.M., and W. Schaffner: "CpG methylation of the cAMP-responsive enhancer/promoter sequence TGACGTCA abolishes specific factor binding as well as transcriptional activation". *Genes & Develop.* 3:612, 1989.).
- 30
- 35

The transcriptional activity of the CRE is increased during B cell activation (Xie, H. T.C. Chiles, and T.L. Rothstein: "Induction of CREB activity via the surface Ig receptor of B cells". *J. Immunol.* 151:880, 1993.). CREB/ATF proteins appear to regulate the expression of multiple genes through the CRE including immunologically important genes such as fos, jun B, Rb-1, IL-6, IL-1 (Tsukada, J., K. Saito, W.R. Waterman, A.C. Webb, and P.E. Auron: "Transcription factors NF-IL6 and CREB recognize a common essential site in the human prointerleukin 1 β gene". *Mol. Cell. Biol.* 14:7285, 1994; Gray, G.D., O.M. Hernandez, D. Hebel, M. Root, J.M. Pow-Sang, and E. Wickstrom: "Antisense DNA inhibition of tumor growth induced by c-Ha-ras oncogene in nude mice". *Cancer Res.* 53:577, 1993), IFN- β (Du, W., and T. Maniatis: "An ATF/CREB binding site protein is required for virus induction of the human interferon B gene". *Proc. Natl. Acad. Sci. USA* 89:2150, 1992), TGF- β 1 (Asiedu, C.K., L. Scott, R.K. Assoian, M. Ehrlich: "Binding of AP-1/CREB proteins and of MDBP to contiguous sites downstream of the human TGF-B1 gene". *Biochim. Biophys. Acta* 1219:55, 1994.), TGF- β 2, class II MHC (Cox, P.M., and C.R. Goding: "An ATF/CREB binding motif is required for aberrant constitutive expression of the MHC class II DR α promoter and activation by SV40 T-antigen". *Nucl. Acids Res.* 20:4881, 1992.), E-selectin, GM-CSF, CD-8 α , the germline Ig α constant region gene, the TCR V β gene, and the proliferating cell nuclear antigen (Huang, D., P.M. Shipman-Appasamy, D.J. Orten, S.H. Hinrichs, and M.B. Prystowsky: "Promoter activity of the proliferating-cell nuclear antigen gene is associated with inducible CRE-binding proteins in interleukin 2-stimulated T lymphocytes". *Mol. Cell. Biol.* 14:4233, 1994.). In addition to activation through the cAMP pathway, CREB can also mediate transcriptional responses to changes in intracellular Ca⁺⁺ concentration (Sheng, M., G. McFadden, and M.E. Greenberg: "Membrane depolarization and calcium induce c-fos transcription via phosphorylation of transcription factor CREB". *Neuron* 4:571, 1990).

The role of protein-protein interactions in transcriptional activation by CREB/ATF proteins appears to be extremely important. Activation of CREB through the cyclic AMP pathway requires protein kinase A (PKA), which phosphorylates CREB³⁴¹ on ser¹³³ and allows it to bind to a recently cloned protein, CBP (Kwok, R.P.S., J.R. Lundblad, J.C. Chrivia, J.P. Richards, H.P. Bachinger, R.G. Brennan, S.G.E. Roberts, M.R. Green, and R.H. Goodman: "Nuclear protein CBP is a coactivator for the transcription factor CREB". *Nature* 370:223, 1994; Arias, J., A.S. Alberts, P. Brindle, F.X. Claret, T. Smea, M. Karin, J. Feramisco, and M. Montminy: "Activation of cAMP and mitogen responsive genes relies on a common nuclear factor". *Nature* 370:226, 1994.). CBP in turn interacts with the basal transcription factor TFIIB causing increased transcription. CREB also has been reported to interact with dTAFII 110, a TATA binding protein-associated factor whose binding may

regulate transcription (Ferrer, K., G. Gill, and M. Montminy: "The cAMP-regulated transcription factor CREB interacts with a component of the TFIID complex". *Proc. Natl. Acad. Sci. USA* 91:1210, 1994.). In addition to these interactions, CREB/ATF proteins can specifically bind multiple other nuclear factors (Hoeffler, J.P., J.W. Lustbader, and C.-Y. Chen: "Identification of multiple nuclear factors that interact with cyclic adenosine 3',5'-monophosphate response element-binding protein and activating transcription factor-2 by protein-protein interactions". *Mol. Endocrinol.* 5:256, 1991) but the biologic significance of most of these interactions is unknown. CREB is normally thought to bind DNA either as a homodimer or as a heterodimer with several other proteins. Surprisingly, CREB monomers constitutively activate transcription (Krajewski, W., and K.A.W. Lee: "A monomeric derivative of the cellular transcription factor CREB functions as a constitutive activator". *Mol. Cell. Biol.* 14:7204, 1994.).

Aside from their critical role in regulating cellular transcription, it has recently been shown that CREB/ATF proteins are subverted by some infectious viruses and retroviruses, which require them for viral replication. For example, the cytomegalovirus immediate early promoter, one of the strongest known mammalian promoters, contains eleven copies of the CRE which are essential for promoter function (Chang, Y.-N., S. Crawford, J. Stall, D.R. Rawlins, K.-T. Jeang, and G.S. Hayward: "The palindromic series I repeats in the simian cytomegalovirus major immediate-early promoter behave as both strong basal enhancers and cyclic AMP response elements". *J. Virol.* 64:264, 1990). At least some of the transcriptional activating effects of the adenovirus E1A protein, which induces many promoters, are due to its binding to the DNA binding domain of the CREB/ATF protein, ATF-2, which mediates E1A inducible transcription activation (Liu, F., and M.R. Green: "Promoter targeting by adenovirus E1a through interaction with different cellular DNA-binding domains". *Nature* 368:520, 1994). It has also been suggested that E1A binds to the CREB-binding protein, CBP (Arany, Z., W.R. Sellers, D.M. Livingston, and R. Eckner: "E1A-associated p300 and CREB-associated CBP belong to a conserved family of coactivators". *Cell* 77:799, 1994). Human T lymphotropic virus-I (HTLV-1), the retrovirus which causes human T cell leukemia and tropical spastic paresis, also requires CREB/ATF proteins for replication. In this case, the retrovirus produces a protein, Tax, which binds to CREB/ATF proteins and redirects them from their normal cellular binding sites to different DNA sequences (flanked by G- and C-rich sequences) present within the HTLV transcriptional enhancer (Paca-Uccaralertkun, S., L.-J. Zhao, N. Adya, J.V. Cross, B.R. Cullen, I.M. Boros, and C.-Z. Giam: "In vitro selection of DNA elements highly responsive to the human T-cell lymphotropic virus type I transcriptional activator, Tax". *Mol. Cell. Biol.* 14:456, 1994; Adya, N., L.-J. Zhao, W. Huang, I. Boros, and C.-Z. Giam: "Expansion of CREB's DNA recognition specificity by Tax results from interaction with Ala-Ala-Arg at

positions 282-284 near the conserved DNA-binding domain of CREB". *Proc. Natl. Acad. Sci. USA* 91:5642, 1994).

5 Summary of the Invention

The instant invention is based on the finding that certain oligonucleotides containing unmethylated cytosine-guanine (CpG) dinucleotides activate lymphocytes as evidenced by *in vitro* and *in vivo* data. Based on this finding, the invention features, in one aspect, novel immunostimulatory oligonucleotide compositions.

In a preferred embodiment, an *immunostimulatory oligonucleotide* is synthetic, between 2 to 100 base pairs in size and contains a consensus mitogenic CpG motif represented by the formula:



wherein C and G are unmethylated, X_1 , X_2 , X_3 and X_4 are nucleotides and a GCG trinucleotide sequence is not present at or near the 5' and 3' termini.

For facilitating uptake into cells, CpG containing immunostimulatory oligonucleotides are preferably in the range of 8 to 40 base pairs in size. Prolonged immunostimulation can be obtained using stabilized oligonucleotides, particularly phosphorothioate stabilized oligonucleotides. Enhanced immunostimulatory activity has been observed where X_1X_2 is the dinucleotide GpA and/or X_3X_4 is the dinucleotide is most preferably TpC or also TpT. Further enhanced immunostimulatory activity has been observed where the consensus motif $X_1X_2CGX_3X_4$ is preceded on the 5' end by a T.

In a second aspect, the invention features useful methods, which are based on the immunostimulatory activity of the oligonucleotides. For example, lymphocytes can either be obtained from a subject and stimulated *ex vivo* upon contact with an appropriate oligonucleotide; or a non-methylated CpG containing oligonucleotide can be administered to a subject to facilitate *in vivo* activation of a subject's lymphocytes. Activated lymphocytes, stimulated by the methods described herein (e.g. either *ex vivo* or *in vivo*), can boost a subject's immune response. The immunostimulatory oligonucleotides can therefore be used to treat, prevent or ameliorate an immune system deficiency (e.g., a tumor or cancer or a viral, fungal, bacterial or parasitic infection in a subject. In addition, immunostimulatory oligonucleotides can also be administered as a vaccine adjuvant, to stimulate a subject's

- 8 -

response to a vaccine. Further, the ability of immunostimulatory cells to induce leukemic cells to enter the cell cycle, suggests a utility for treating leukemia by increasing the sensitivity of chronic leukemia cells and then administering conventional ablative chemotherapy.

In a third aspect, the invention features *neutral oligonucleotides* (i.e. oligonucleotide that do not contain an unmethylated CpG or which contain a methylated CpG dinucleotide). In a preferred embodiment, a neutralizing oligonucleotide is complementary to an immunostimulatory sequence, but contains a methylated instead of an unmethylated CpG dinucleotide sequence and therefore can compete for binding with unmethylated CpG containing oligonucleotides. In a preferred embodiment, the methylation occurs at one or more of the four carbons and two nitrogens comprising the cytosine six member ring or at one or more of the five carbons and four nitrogens comprising the guanine nine member double ring. 5' methyl cytosine is a preferred methylated CpG.

In a fourth aspect, the invention features useful methods using the neutral oligonucleotides. For example, in vivo administration of neutral oligonucleotides should prove useful for treating diseases such as systemic lupus erythematosus, sepsis and autoimmune diseases, which are caused or exacerbated by the presence of unmethylated CpG dimers in a subject. In addition, methylation CpG containing antisense oligonucleotides or oligonucleotide probes would not initiate an immune reaction when administered to a subject *in vivo* and therefore would be safer than corresponding unmethylated oligonucleotides.

In a fifth aspect, the invention features *immunoinhibitory oligonucleotides*, which are capable of interfering with the activity of viral or cellular transcription factors. In a preferred embodiment, immunoinhibitory oligonucleotides are between 2 to 100 base pairs in size and contain a consensus immunoinhibitory CpG motif represented by the formula:



wherein X = a nucleotide and n = in the range of 0-50. In a preferred embodiment, X is a pyrimidine.

For facilitating uptake into cells, immunoinhibitory oligonucleotides are preferably in the range of 8 to 40 base pairs in size. Prolonged biologic effects can be obtained using stabilized oligonucleotides, particularly phosphorothioate stabilized oligonucleotides.

In a sixth and final aspect, the invention features various uses for immunoinhibitory oligonucleotides. Immunoinhibitory oligonucleotides have antiviral activity, independent of any antisense effect due to complementarity between the oligonucleotide and the viral sequence being targeted.

5

Other features and advantages of the invention will become more apparent from the following detailed description and claims.

10 Detailed Description of the Invention

Definitions

As used herein, the following terms and phrases shall have the meanings set forth below:

15

An "oligonucleotide" or "oligo" shall mean multiple nucleotides (i.e. molecules comprising a sugar (e.g. ribose or deoxyribose) linked to a phosphate group and to an exchangeable organic base, which is either a substituted pyrimidine (e.g. cytosine (C), thymine (T) or uracil (U)) or a substituted purine (e.g. adenine (A) or guanine (G)). The term "oligonucleotide" as used herein refers to both oligoribonucleotides (ORNs) and oligodeoxyribonucleotides (ODNs). The term "oligonucleotide" shall also include oligonucleosides (i.e. an oligonucleotide minus the phosphate) and any other organic base containing polymer. Oligonucleotides can be obtained from existing nucleic acid sources (e.g. genomic or cDNA), but are preferably synthetic (e.g. produced by oligonucleotide synthesis).

25

A "stabilized oligonucleotide" shall mean an oligonucleotide that is relatively resistant to in vivo degradation (e.g. via an exo- or endo-nuclease). Preferred stabilized oligonucleotides of the instant invention have a modified phosphate backbone. Especially preferred oligonucleotides have a phosphorothioate modified phosphate backbone (i.e. at least one of the phosphate oxygens is replaced by sulfur). Other stabilized oligonucleotides include: nonionic DNA analogs, such as alkyl- and aryl- phosphonates (in which the charged phosphonate oxygen is replaced by an alkyl or aryl group), phosphodiester and alkylphosphotriesters, in which the charged oxygen moiety is alkylated. Oligonucleotides which contain a diol, such as tetraethyleneglycol or hexaethyleneglycol, at either or both termini have also been shown to be substantially resistant to nuclease degradation.

35

- 10 -

An "immunostimulatory oligonucleotide", "immunostimulatory CpG containing oligonucleotide", or "CpG ODN" refer to an oligonucleotide, which contains a cytosine, guanine dinucleotide sequence and stimulates (e.g. has a mitogenic effect) on vertebrate lymphocyte. Preferred immunostimulatory oligonucleotides are between 2 to 100 base pairs in size and contain a consensus mitogenic CpG motif represented by the formula:



wherein C and G are unmethylated, X_1 , X_2 , X_3 and X_4 are nucleotides and a GCG trinucleotide sequence is not present at or near the 5' and 3' termini.

Preferably the immunostimulatory oligonucleotides range between 8 to 40 base pairs in size. In addition, the immunostimulatory oligonucleotides are preferably stabilized oligonucleotides, particularly preferred are phosphorothioate stabilized oligonucleotides. In one preferred embodiment, X_1X_2 is the dinucleotide GpA. In another preferred embodiment, X_3X_4 is preferably the dinucleotide TpC or also TpT. In a particularly preferred embodiment, the consensus motif $X_1X_2CGX_3X_4$ is preceded on the 5' end by a T. Particularly preferred consensus sequences are TGACGTT or TGACGTC.

A "neutral oligonucleotide" refers to an oligonucleotide that does not contain an unmethylated CpG or an oligonucleotide which contains a methylated CpG dinucleotide. In a preferred embodiment, a neutralizing oligonucleotide is complementary to an immunostimulatory sequence, but contains a methylated instead of an unmethylated CpG dinucleotide sequence and therefore can compete for binding with unmethylated CpG containing oligonucleotides. In a preferred embodiment, the methylation occurs at one or more of the four carbons and two nitrogens comprising the cytosine six member ring or at one or more of the five carbons and four nitrogens comprising the guanine nine member double ring. 5' methyl cytosine is a preferred methylated CpG.

An "immunoinhibitory oligonucleotide" or "immunoinhibitory CpG containing oligonucleotide" is an oligonucleotide that is capable of interfering with the activity of viral or cellular transcription factors (support in summary of the invention, page 8). Preferable immunoinhibitory oligonucleotides are between 2 to 100 base pairs in size and can be represented by the formula:



wherein X = a nucleotide and n = in the range of 0-50. In a preferred embodiment, X is a pyrimidine.

- 11 -

For facilitating uptake into cells, immunoinhibitory oligonucleotides are preferably in the range of 8 to 40 base pairs in size. Prolonged immunostimulation can be obtained using stabilized oligonucleotides, particularly phosphorothioate stabilized.

"Palindromic sequence" shall mean an inverted repeat (i.e. a sequence such as ABCDEE'D'C'B'A' in which A and A' are bases capable of forming the usual Watson-Crick base pairs. In vivo, such sequences may form double stranded structures.

An "oligonucleotide delivery complex" shall mean an oligonucleotide associated with (e.g. ionically or covalently bound to; or encapsulated within) a targeting means (e.g. a molecule that results in higher affinity binding to target cell (e.g. B-cell and natural killer (NK) cell) surfaces and/or increased cellular uptake by target cells). Examples of oligonucleotide delivery complexes include oligonucleotides associated with: a sterol (e.g. cholesterol), a lipid (e.g. a cationic lipid, virosome or liposome), or a target cell specific binding agent (e.g. a ligand recognized by target cell specific receptor). Preferred complexes must be sufficiently stable in vivo to prevent significant uncoupling prior to internalization by the target cell. However, the complex should be cleavable under appropriate conditions within the cell so that the oligonucleotide is released in a functional form.

An "immune system deficiency" shall mean a disease or disorder in which the subject's immune system is not functioning in normal capacity or in which it would be useful to boost a subject's immune response for example to eliminate a tumor or cancer (e.g. tumors of the brain, lung (e.g. small cell and non-small cell), ovary, breast, prostate, colon, as well as other carcinomas and sarcomas) or a viral (e.g. HIV, herpes), fungal (e.g. *Candida sp.*), bacterial or parasitic (e.g. *Leishmania*, *Toxoplasma*) infection in a subject.

A "disease associated with immune system activation" shall mean a disease or condition caused or exacerbated by activation of the subject's immune system. Examples include systemic lupus erythematosus, sepsis and autoimmune diseases such as rheumatoid arthritis and multiple sclerosis.

A "subject" shall mean a human or vertebrate animal including a dog, cat, horse, cow, pig, sheep, goat, chicken, monkey, rat, mouse, etc.

Certain Unmethylated CpG Containing Oligos Have B Cell Stimulatory Activity As Shown in vitro and in vivo

5 In the course of investigating the lymphocyte stimulatory effects of two antisense oligonucleotides specific for endogenous retroviral sequences, using protocols described in the attached Examples 1 and 2, it was surprisingly found that two out of twenty-four "controls" (including various scrambled, sense, and mismatch controls for a panel of "antisense" ODN) also mediated B cell activation and IgM secretion, while the other "controls" had no effect.

10

Two observations suggested that the mechanism of this B cell activation by the "control" ODN may not involve antisense effects 1) comparison of vertebrate DNA sequences listed in GenBank showed no greater homology than that seen with non-stimulatory ODN and 2) the two controls showed no hybridization to Northern blots with 10 μ g of spleen poly A+ RNA. Resynthesis of these ODN on a different synthesizer or extensive purification by polyacrylamide gel electrophoresis or high pressure liquid chromatography gave identical stimulation, eliminating the possibility of an impurity. 15 Similar stimulation was seen using B cells from C3H/HeJ mice, eliminating the possibility that lipopolysaccharide (LPS) contamination could account for the results.

20

The fact that two "control" ODN caused B cell activation similar to that of the two "antisense" ODN raised the possibility that all four ODN were stimulating B cells through some non-antisense mechanism involving a sequence motif that was absent in all of the other nonstimulatory control ODN. In comparing these sequences, it was discovered that 25 all of the four stimulatory ODN contained ODN dinucleotides that were in a different sequence context from the nonstimulatory control.

To determine whether the CpG motif present in the stimulatory ODN was responsible for the observed stimulation, over 300 ODN ranging in length from 5 to 42 bases 30 that contained methylated, unmethylated, or no CpG dinucleotides in various sequence contexts were synthesized. These ODNs, including the two original "controls" (ODN 1 and 2) and two originally synthesized as "antisense" (ODN 3D and 3M; Krieg, A.M. *J. Immunol.* 143:2448 (1989)), were then examined for in vitro effects on spleen cells (representative sequences are listed in Table 1). Several ODN that contained CpG dinucleotides induced B 35 cell activation and IgM secretion; the magnitude of this stimulation typically could be increased by adding more CpG dinucleotides (Table 1; compare ODN 2 to 2a or 3D to 3Da and 3Db). Stimulation did not appear to result from an antisense mechanism or impurity. ODN caused no detectable activation of $\gamma\delta$ or other T cell populations.

Mitogenic ODN sequences uniformly became nonstimulatory if the CpG dinucleotide was mutated (Table 1; compare ODN 1 to 1a; 3D to 3Dc; 3M to 3Ma; and 4 to 4a) or if the cytosine of the CpG dinucleotide was replaced by 5-methylcytosine (Table 1; ODN 1b,2b,2c,3Dd, and 3Mb). In contrast, methylation of other cytosines did not reduce ODN activity (ODN 1c, 2d, 3De and 3Mc). These data confirmed that a CpG motif is the essential element present in ODN that activate B cells.

In the course of these studies, it became clear that the bases flanking the CpG dinucleotide played an important role in determining the B cell activation induced by an ODN. The optimal stimulatory motif was determined to consist of a CpG flanked by two 5' purines (preferably a GpA dinucleotide) and two 3' pyrimidines (preferably a TpT or TpC dinucleotide). Mutations of ODN to bring the CpG motif closer to this ideal improved stimulation (e.g. compare ODN 2 to 2e; 3M to 3Md) while mutations that disturbed the motif reduced stimulation (e.g. compare ODN 3D to 3Df; 4 to 4b, 4c and 4d). On the other hand, mutations outside the CpG motif did not reduce stimulation (e.g. compare ODN 1 to 1d; 3D to 3Dg; 3M to 3Me).

Of those tested, ODNs shorter than 8 bases were non-stimulatory (e.g. ODN 4e). Among the forty-eight 8 base ODN tested, the most stimulatory sequence identified was TCAACGTT (ODN 4) which contains the self complementary "palindrome" AACGTT. In further optimizing this motif, it was found that ODN containing Gs at both ends showed increased stimulation, particularly if the the ODN were rendered nuclease resistant by phosphorothioate modification of the terminal internucleotide linkages. ODN 1585 (5' GGGGTCAACGTTTCAGGGGGG 3' (SEQ ID NO:1)), in which the first two and last five internucleotide linkages are phosphorothioate modified caused an average 25.4 fold increase in mouse spleen cell proliferation compared to an average 3.2 fold increase in proliferation induced by ODN 1638, which has the same sequence as ODN 1585 except that the 10 Gs at the two ends are replaced by 10 As. The effect of the G-rich ends is *cis*; addition of an ODN with poly G ends but no CpG motif to cells along with 1638 gave no increased proliferation.

Other octamer ODN containing a 6 base palindrome with a TpC dinucleotide at the 5' end were also active if they were close to the optimal motif (e.g. ODN 4b,4c). Other dinucleotides at the 5' end gave reduced stimulation (eg ODN 4f; all sixteen possible dinucleotides were tested). The presence of a 3' dinucleotide was insufficient to compensate for the lack of a 5' dinucleotide (eg. ODN 4g). Disruption of the palindrome eliminated stimulation in octamer ODN (eg., ODN 4h), but palindromes were not required in longer ODN.

Table 1: Oligonucleotide Stimulation of B Cells

5

10

ODN	Sequence (5' to 3')†	Stimulation Index'	
		³ H Uridine	IgM Production
1 (SEQ ID NO: 2)	GCTAGACGTTAGCGT	6.1 ± 0.8	17.9 ± 3.6
1a (SEQ ID NO: 3)T.....	1.2 ± 0.2	1.7 ± 0.5
1b (SEQ ID NO: 4)Z.....	1.2 ± 0.1	1.8 ± 0.0
1c (SEQ ID NO: 5)Z..	10.3 ± 4.4	9.5 ± 1.8
1d (SEQ ID NO: 6)	..AT.....GAGC.	13.0 ± 2.3	18.3 ± 7.5
2 (SEQ ID NO: 7)	ATGGAAGGTCCAGCGTTCTC	2.9 ± 0.2	13.6 ± 2.0
2a (SEQ ID NO: 8)	..C..CTC..G.....	7.7 ± 0.8	24.2 ± 3.2
2b (SEQ ID NO: 9)	..Z..CTC..ZG..Z.....	1.6 ± 0.5	2.8 ± 2.2
2c (SEQ ID NO: 10)	..Z..CTC..G.....	3.1 ± 0.6	7.3 ± 1.4
2d (SEQ ID NO: 11)	..C..CTC..G.....Z..	7.4 ± 1.4	27.7 ± 5.4
2e (SEQ ID NO: 12)A.....	5.6 ± 2.0	ND
3D (SEQ ID NO: 13)	GAGAAAGCTGGACCTTCCAT	4.9 ± 0.5	19.9 ± 3.6
3Da (SEQ ID NO: 14)C.....	6.6 ± 1.5	33.9 ± 6.8
3Db (SEQ ID NO: 15)C.....G..	10.1 ± 2.8	25.4 ± 0.8
3Dc (SEQ ID NO: 16)	...C.A.....	1.0 ± 0.1	1.2 ± 0.5
3Dd (SEQ ID NO: 17)Z.....	1.2 ± 0.2	1.0 ± 0.4
3De (SEQ ID NO: 18)Z.....	4.4 ± 1.2	18.8 ± 4.4
3Df (SEQ ID NO: 19)A.....	1.6 ± 0.1	7.7 ± 0.4
3Dg (SEQ ID NO: 20)CC.G.ACTG..	6.1 ± 1.5	18.6 ± 1.5
3M (SEQ ID NO: 21)	TCCATGTGGTCTTGATGCT	4.1 ± 0.2	23.2 ± 4.9
3Ma (SEQ ID NO: 22)CT.....	0.9 ± 0.1	1.8 ± 0.5
3Mb (SEQ ID NO: 23)Z.....	1.3 ± 0.3	1.5 ± 0.6
3Mc (SEQ ID NO: 24)Z.....	5.4 ± 1.5	8.5 ± 2.6
3Md (SEQ ID NO: 25)AT.....	17.2 ± 9.4	ND
3Me (SEQ ID NO: 26)C..A..	3.6 ± 0.2	14.2 ± 5.2

- 15 -

4	TCAACGTT	6.1 ± 1.4	19.2 ± 5.2
4aGC..	1.1 ± 0.2	1.5 ± 1.1
4b	...GCGC.	4.5 ± 0.2	9.6 ± 3.4
4c	...TCGA.	2.7 ± 1.0	ND
4d	..TT...AA	1.3 ± 0.2	ND
4e	-...-...-	1.3 ± 0.2	1.1 ± 0.5
4f	C...-...-	3.9 ± 1.4	ND
4g	-...-...CT	1.4 ± 0.3	ND
4h-...C	1.2 ± 0.2	ND
LPS		7.8 ± 2.5	4.8 ± 1.0

' Stimulation indexes are the means and std. dev. derived from at least 3 separate experiments, and are compared to wells cultured with no added ODN. ND = not done. CpG dinucleotides are underlined. Dots indicate identity; dashes indicate deletions. Z indicates 5 methyl cytosine.)

5

The kinetics of lymphocyte activation were investigated using mouse spleen cells. When the cells were pulsed at the same time as ODN addition and harvested just four hours later, there was already a two-fold increase in ³H uridine incorporation. Stimulation peaked at 12-48 hours and then decreased. After 24 hours, no intact ODN were detected, perhaps accounting for the subsequent fall in stimulation when purified B cells with or without anti-IgM (at a submitogenic dose) were cultured with CpG ODN, proliferation was found to synergistically increase about 10-fold by the two mitogens in combination after 48 hours. The magnitude of stimulation was concentration dependent and consistently exceeded that of LPS under optimal conditions for both. Oligonucleotides containing a nuclease resistant phosphorothioate backbone were approximately two hundred times more potent than unmodified oligonucleotides.

Cell cycle analysis was used to determine the proportion of B cells activated by CpG-ODN. CpG-ODN induced cycling in more than 95% of B cells (Table 2). Splenic B lymphocytes sorted by flow cytometry into CD23- (marginal zone) and CD23+ (follicular) subpopulations were equally responsive to ODN- induced stimulation, as were both resting and activated populations of B cells isolated by fractionation over Percoll gradients. These studies demonstrated that CpG-ODN induce essentially all B cells to enter the cell cycle.

SUBSTITUTE SHEET (RULE 26)

Table 2: Cell Cycle Analysis with CpG ODN

Treatment	Percent of cells in		
	G0	G1	SA+G2+M
Media	97.6	2.4	0.02
ODN 1a	95.2	4.8	0.04
ODN 1d	2.7	74.4	22.9
ODN 3Db	3.5	76.4	20.1
LPS (30 µg/ml)	17.3	70.5	12.2

The mitogenic effects of CpG ODN on human cells, were tested on peripheral blood mononuclear cells (PBMCs) obtained from two patients with chronic lymphocytic leukemia (CLL), as described in Example 1. Control ODN containing no CpG dinucleotide sequence showed no effect on the basal proliferation of 442 cpm and 874 cpm (proliferation measured by ³H thymidine incorporation) of the human cells. However, a phosphorothioate modified CpG ODN 3Md (SEQ ID NO: 25) induced increased proliferation of 7210 and 86795 cpm respectively in the two patients at a concentration of just 1 μM. Since these cells had been frozen, they may have been less responsive to the oligos than fresh cells *in vivo*. In addition, cells from CLL patients typically are non-proliferating, which is why traditional chemotherapy is not effective.

Certain B cell lines such as WEHI-231 are induced to undergo growth arrest and/or apoptosis in response to crosslinking of their antigen receptor by anti-IgM (Jakway, J.P. et al., "Growth regulation of the B lymphoma cell line WEHI-231 by anti-immunoglobulin, lipopolysaccharide and other bacterial products" *J. Immunol.* 137: 2225 (1986); Tsubata, T., J. Wu and T. Honjo: B-cell apoptosis induced by antigen receptor crosslinking is blocked by a T-cell signal through CD40." *Nature* 364: 645 (1993)). WEHI-231 cells are rescued from this growth arrest by certain stimuli such as LPS and by the CD40 ligand. ODN containing the CpG motif were also found to protect WEHI-231 from anti-IgM induced growth arrest, indicating that accessory cell populations are not required for the effect.

To better understand the immune effects of unmethylated CpG ODN, the levels of cytokines and prostaglandins *in vitro* and *in vivo* were measured. Unlike LPS, CpG

ODN were not found to induce purified macrophages to produce prostaglandin PGE₂. In fact, no apparent direct effect of CpG ODN was detected on either macrophages or T cells. *In vivo* or in whole spleen cells, no significant increase in the following interleukins: IL-2, IL-3, IL-4, or IL-10 was detected within the first six hours. However, the level of IL-6 increased strikingly within 2 hours in the serum of mice injected with CpG ODN. Increased expression of IL-12 and interferon gamma (IFN- γ) by spleen cells was also detected within the first two hours.

To determine whether CpG ODN can cause *in vivo* immune stimulation, DBA/2 mice were injected once intraperitoneally with PBS or phosphorothioate CpG or non-CpG ODN at a dose of 33 mg/kg (approximately 500 μ g/mouse). Pharmacokinetic studies in mice indicate that this dose of phosphorothioate gives levels of approximately 10 μ g/g in spleen tissue (within the effective concentration range determined from the *in vitro* studies described herein) for longer than twenty-four hours (Agrawal, S. et al. (1991) *Proc. Natl. Acad. Sci. USA* 91:7595). Spleen cells from mice were examined twenty-four hours after ODN injection for expression of B cells surface activation markers Ly-6A/E, B220, and class II MHC using three color flow cytometry and for their spontaneous proliferation using ³H thymidine. Expression of all three activation markers was significantly increased in B cells from mice injected with CpG ODN, but not from mice injected with PBS or non-CpG ODN. Spontaneous ³H thymidine incorporation was increased by 2-6 fold in spleen cells from mice injected with the stimulatory ODN compared to PBS or non-CpG ODN-injected mice. After 4 days, serum IgM levels in mice injected with CpG ODN *in vivo* were increased by approximately 3-fold compared to controls. Consistent with the inability of these agents to activate T cells, there was minimal change in T cell expression of the IL-2R or CD-44.

Degradation of phosphodiester ODN in serum is predominantly mediated by 3' exonucleases, while intracellular ODN degradation is more complex, involving 5' and 3' exonucleases and endonucleases. Using a panel of ODN bearing the 3D sequence with varying numbers of phosphorothioate modified linkages at the 5' and 3' ends, it was empirically determined that two 5' and five 3' modified linkages are required to provide optimal stimulation with this CpG ODN.

Unmethylated CpG Containing Oligos Have NK Cell Stimulatory Activity

As described in further detail in Example 4, experiments were conducted to determine whether CpG containing oligonucleotides stimulated the activity of natural killer (NK) cells in addition to B cells. As shown in Table 3, a marked induction of NK activity among spleen cells cultured with CpG ODN 1 and 3Dd was observed. In contrast, there was relatively no induction in effectors that had been treated with non-CpG control ODN.

Table 3: Induction Of NK Activity By CpG Oligodeoxynucleotides (ODN)

ODN	% YAC-1 Specific Lysis*		% 2C11 Specific Lysis	
	Effector: Target		Effector: Target	
	50:1	100:1	50:1	100:1
None	-1.1	-1.4	15.3	16.6
1	16.1	24.5	38.7	47.2
3Dd	17.1	27.0	37.0	40.0
non-CpG ODN	-1.6	-1.7	14.8	15.4

5

Neutralizing Activity of Methylated CpG Containing Oligos

10 B cell mitogenicity of ODN in which cytosines in CpG motifs or elsewhere were replaced by 5-methylcytosine were tested as described in Example 1. As shown in Table 1 above, ODN containing methylated CpG motifs were non-mitogenic (Table 1; ODN 1c, 2f, 3De, and 3Mc). However, methylation of cytosines other than in a CpG dinucleotide retained their stimulatory properties (Table 1, ODN 1d, 2d, 3Df, and 3Md).

15

Immunoinhibitory Activity of Oligos Containing a GCG Trinucleotide Sequence at or near both termini

20 In some cases, ODN containing CpG dinucleotides that are not in the stimulatory motif described above were found to block the stimulatory effect of other mitogenic CpG ODN. Specifically the addition of an atypical CpG motif consisting of a GCG near or at the 5' and/or 3' end of CpG ODN actually inhibited stimulation of proliferation by other CpG motifs. Methylation or substitution of the cytosine in a GCG motif reverses this effect. By itself, a GCG motif in an ODN has a modest mitogenic effect, though far lower
25 than that seen with the preferred CpG motif.

Proposed Mechanisms of Action of Immunostimulatory, Neutralizing and Immunoinhibitory Oligonucleotides

30 Unlike antigens that trigger B cells through their surface Ig receptor, CpG-ODN did not induce any detectable Ca^{2+} flux, changes in protein tyrosine phosphorylation, or IP 3 generation. Flow cytometry with FITC-conjugated ODN with or without a CpG motif was performed as described in Zhao, Q et al., (*Antisense Research and Development* 3:53-66

(1993)), and showed equivalent membrane binding, cellular uptake, efflux, and intracellular localization. This suggests that there may not be cell membrane proteins specific for CpG ODN. Rather than acting through the cell membrane, that data suggests that unmethylated CpG containing oligonucleotides require cell uptake for activity: ODN covalently linked to a solid Teflon support were nonstimulatory, as were biotinylated ODN immobilized on either
5 avidin beads or avidin coated petri dishes. CpG ODN conjugated to either FITC or biotin retained full mitogenic properties, indicating no steric hindrance.

The optimal CpG motif (TGACGTT/C is identical to the CRE (cyclic AMP
10 response element). Like the mitogenic effects of CpG ODN, binding of CREB to the CRE is abolished if the central CpG is methylated. Electrophoretic mobility shift assays were used to determine whether CpG ODN, which are single stranded, could compete with the binding of B cell CREB/ATF proteins to their normal binding site, the doublestranded CRE. Competition assays demonstrated that single stranded ODN containing CpG motifs could
15 completely compete the binding of CREB to its binding site, while ODN without CpG motifs could not. These data support the conclusion that CpG ODN exert their mitogenic effects through interacting with one or more B cell CREB/ATF proteins in some way. Conversely, the presence of GCG sequences or other atypical CPG motifs near the 5' and/or 3' ends of ODN likely interact with CREB/ATF proteins in a way that does not cause activation, and
20 may even prevent it.

The stimulatory CpG motif is common in microbial genomic DNA, but quite rare in vertebrate DNA. In addition, bacterial DNA has been reported to induce B cell proliferation and immunoglobulin (Ig) production, while mammalian DNA does not
25 (Messina, J.P. et al., *J. Immunol.* 147:1759 (1991)). Experiments further described in Example 3, in which methylation of bacterial DNA with CpG methylase was found to abolish mitogenicity, demonstrates that the difference in CpG status is the cause of B cell stimulation by bacterial DNA. This data supports the following conclusion: that unmethylated CpG dinucleotides present within bacterial DNA are responsible for the stimulatory effects of
30 bacterial DNA.

Teleologically, it appears likely that lymphocyte activation by the CpG motif represents an immune defense mechanism that can thereby distinguish bacterial from host DNA. Host DNA would induce little or no lymphocyte activation due to it CpG suppression
35 and methylation. Bacterial DNA would cause selective lymphocyte activation in infected tissues. Since the CpG pathway synergizes with B cell activation through the antigen receptor, B cells bearing antigen receptor specific for bacterial antigens would receive one activation signal through cell membrane Ig and a second signal from bacterial DNA, and

would therefore tend to be preferentially activated. The interrelationship of this pathway with other pathways of B cell activation provide a physiologic mechanism employing a polyclonal antigen to induce antigen-specific responses.

5

Method for Making Immunostimulatory Oligos

For use in the instant invention, oligonucleotides can be synthesized de novo using any of a number of procedures well known in the art. For example, the β -cyanoethyl phosphoramidite method (S.L. Beaucage and M.H. Caruthers, (1981) *Tet. Let.* 22:1859);
10 nucleoside H-phosphonate method (Garegg et al., (1986) *Tet. Let.* 27: 4051-4054; Froehler et al., (1986) *Nucl. Acid. Res.* 14: 5399-5407; Garegg et al., (1986) *Tet. Let.* 27: 4055-4058, Gaffney et al., (1988) *Tet. Let.* 29:2619-2622). These chemistries can be performed by a variety of automated oligonucleotide synthesizers available in the market. Alternatively, oligonucleotides can be prepared from existing nucleic acid sequences (e.g. genomic or
15 cDNA) using known techniques, such as those employing restriction enzymes, exonucleases or endonucleases.

For use *in vivo*, oligonucleotides are preferably relatively resistant to degradation (e.g. via endo- and exo- nucleases). Oligonucleotide stabilization can be
20 accomplished via phosphate backbone modifications. A preferred stabilized oligonucleotide has a phosphorothioate modified backbone. The pharmacokinetics of phosphorothioate ODN show that they have a systemic half-life of forty-eight hours in rodents and suggest that they may be useful for *in vivo* applications (Agrawal, S. et al. (1991) *Proc. Natl. Acad. Sci. USA* 88:7595). Phosphorothioates may be synthesized using automated techniques employing
25 either phosphoramidate or H phosphonate chemistries. Aryl- and alkyl- phosphonates can be made e.g. (as described in U.S. Patent No. 4,469,863); and alkylphosphotriesters (in which the charged oxygen moiety is alkylated as described in U.S. Patent No. 5,023,243 and European Patent No. 092,574) can be prepared by automated solid phase synthesis using commercially available reagents. Methods for making other DNA backbone modifications
30 and substitutions have been described (Uhlmann, E. and Peyman, A. (1990) *Chem. Rev.* 90:544; Goodchild, J. (1990) *Bioconjugate Chem.* 1:165).

For administration *in vivo*, oligonucleotides may be associated with a molecule that results in higher affinity binding to target cell (e.g. B-cell and natural killer
35 (NK) cell) surfaces and/or increased cellular uptake by target cells to form an "olig nucleotide delivery complex". Oligonucleotides can be ionically, or covalently associated with appropriate molecules using techniques which are well known in the art. A variety of coupling or crosslinking agents can be used e.g. protein A, carbodiimide, and N-

succinimidyl-3-(2-pyridyldithi) propionate (SPDP). Oligonucleotides can alternatively be encapsulated in liposomes or virosomes using well-known techniques.

The present invention is further illustrated by the following Examples which in no way should be construed as further limiting. The entire contents of all of the references (including literature references, issued patents, published patent applications, and co-pending patent applications) cited throughout this application are hereby expressly incorporated by reference.

Therapeutic Uses of Immunostimulatory Oligos

Based on their immunostimulatory properties, oligonucleotides containing at least one unmethylated CpG dinucleotide can be administered to a subject in vivo to treat an "immune system deficiency". Alternatively, oligonucleotides containing at least one unmethylated CpG dinucleotide can be contacted with lymphocytes (e.g. B cells or NK cells) obtained from a subject having an immune system deficiency ex vivo and activated lymphocytes can then be reimplanted in the subject.

Immunostimulatory oligonucleotides can also be administered to a subject in conjunction with a vaccine, as an adjuvant, to boost a subject's immune system to effect better response from the vaccine. Preferably the unmethylated CpG dinucleotide is administered slightly before or at the same time as the vaccine.

Preceding chemotherapy with an immunostimulatory oligonucleotide should prove useful for increasing the responsiveness of the malignant cells to subsequent chemotherapy. CpG ODN also increased natural killer cell activity in both human and murine cells. Induction of NK activity may likewise be beneficial in cancer immunotherapy.

Therapeutic Uses for Neutral Oligonucleotides

Oligonucleotides that are complementary to certain target sequences can be synthesized and administered to a subject in vivo. For example, antisense oligonucleotides hybridize to complementary mRNA, thereby preventing expression of a specific target gene. The sequence-specific effects of antisense oligonucleotides have made them useful research tools for the investigation of protein function. Phase I/II human trials of systemic antisense therapy are now underway for acute myelogenous leukemia and HIV.

In addition, oligonucleotide probes (i.e. oligonucleotides with a detectable label) can be administered to a subject to detect the presence of a complementary sequence based on detection of bound label. In vivo administration and detection of oligonucleotide

probes may be useful for diagnosing certain diseases that are caused or exacerbated by certain DNA sequences (e.g. systemic lupus erythematosus, sepsis and autoimmune diseases).

Antisense oligonucleotides or oligonucleotide probes in which any or all CpG dinucleotide is methylated, would not produce an immune reaction when administered to a subject *in vivo* and therefore would be safer than the corresponding non-methylated CpG containing oligonucleotide.

For use in therapy, an effective amount of an appropriate oligonucleotide alone or formulated as an oligonucleotide delivery complex can be administered to a subject by any mode allowing the oligonucleotide to be taken up by the appropriate target cells (e.g. B-cells and NK cells). Preferred routes of administration include oral and transdermal (e.g. via a patch). Examples of other routes of administration include injection (subcutaneous, intravenous, parenteral, intraperitoneal, intrathecal, etc.). The injection can be in a bolus or a continuous infusion.

An oligonucleotide alone or as an oligonucleotide delivery complex can be administered in conjunction with a pharmaceutically acceptable carrier. As used herein, the phrase "pharmaceutically acceptable carrier" is intended to include substances that can be coadministered with an oligonucleotide or an oligonucleotide delivery complex and allows the oligonucleotide to perform its intended function. Examples of such carriers include solutions, solvents, dispersion media, delay agents, emulsions and the like. The use of such media for pharmaceutically active substances are well known in the art. Any other conventional carrier suitable for use with the oligonucleotides falls within the scope of the instant invention.

The language "effective amount" of an oligonucleotide refers to that amount necessary or sufficient to realize a desired biologic effect. For example, an effective amount of an oligonucleotide containing at least one methylated CpG for treating an immune system deficiency could be that amount necessary to eliminate a tumor, cancer, or bacterial, viral or fungal infection. An effective amount for use as a vaccine adjuvant could be that amount useful for boosting a subject's immune response to a vaccine. An "effective amount" of an oligonucleotide lacking a non-methylated CpG for use in treating a disease associated with immune system activation, could be that amount necessary to outcompete non-methylated CpG containing nucleotide sequences. The effective amount for any particular application can vary depending on such factors as the disease or condition being treated, the particular oligonucleotide being administered, the size of the subject, and the severity of the disease or

condition. One of ordinary skill in the art can empirically determine the effective amount of a particular oligonucleotide without necessitating undue experimentation.

5 The studies reported above indicate that unmethylated CpG containing oligonucleotides are directly mitogenic for lymphocytes (e.g. B cells and NK cells). Together with the presence of these sequences in bacterial DNA, these results suggest that the underrepresentation of CpG dinucleotides in animal genomes, and the extensive methylation of cytosines present in such dinucleotides, may be explained by the existence of an immune defense mechanism that can distinguish bacterial from host DNA. Host DNA would
10 commonly be present in many anatomic regions and areas of inflammation due to apoptosis (cell death), but generally induces little or no lymphocyte activation. However, the presence of bacterial DNA containing unmethylated CpG motifs can cause lymphocyte activation precisely in infected anatomic regions, where it is beneficial. This novel activation pathway provides a rapid alternative to T cell dependent antigen specific B cell activation. However,
15 it is likely that B cell activation would not be totally nonspecific. B cells bearing antigen receptors specific for bacterial products could receive one activation signal through cell membrane Ig, and a second from bacterial DNA, thereby more vigorously triggering antigen specific immune responses.

20 As with other immune defense mechanisms, the response to bacterial DNA could have undesirable consequences in some settings. For example, autoimmune responses to self antigens would also tend to be preferentially triggered by bacterial infections, since autoantigens could also provide a second activation signal to autoreactive B cells triggered by bacterial DNA. Indeed the induction of autoimmunity by bacterial infections is a common
25 clinical observance. For example, the autoimmune disease systemic lupus erythematosus, which is: i) characterized by the production of anti-DNA antibodies; ii) induced by drugs which inhibit DNA methyltransferase (Cornacchia, E.J. et al., *J. Clin. Invest.* 92:38 (1993)); and iii) associated with reduced DNA methylation (Richardson, B., L. et al., *Arth. Rheum* 35:647 (1992)), is likely triggered at least in part by activation of DNA-specific B cells
30 through stimulatory signals provided by CpG motifs, as well as by binding of bacterial DNA to antigen receptors.

 Further, sepsis, which is characterized by high morbidity and mortality due to massive and nonspecific activation of the immune system may be initiated by bacterial DNA
35 and other products released from dying bacteria that reach concentrations sufficient to directly activate many lymphocytes.

Lupus, sepsis and other "diseases associated with immune system activation" may be treated, prevented or ameliorated by administering to a subject oligonucleotides lacking an unmethylated CpG dinucleotide (e.g. oligonucleotides that do not include a CpG motif or oligonucleotides in which the CpG motif is methylated) to block the binding of unmethylated CpG containing nucleic acid sequences. Oligonucleotides lacking an unmethylated CpG motif can be administered alone or in conjunction with compositions that block an immune cell's response to other mitogenic bacterial products (e.g. LPS).

The following serves to illustrate mechanistically how oligonucleotides containing an unmethylated CpG dinucleotide can treat, prevent or ameliorate the disease lupus. Lupus is commonly thought to be triggered by bacterial or viral infections. Such infections have been reported to stimulate the production of nonpathogenic antibodies to single stranded DNA. These antibodies likely recognize primarily bacterial sequences including unmethylated CpGs. As disease develops in lupus, the anti-DNA antibodies shift to pathogenic antibodies that are specific for double-stranded DNA. These antibodies would have increased binding for methylated CpG sequences and their production would result from a breakdown of tolerance in lupus. Alternatively, lupus may result when a patient's DNA becomes hypomethylated, thus allowing anti-DNA antibodies specific for unmethylated CpGs to bind to self DNA and trigger more widespread autoimmunity through the process referred to as "epitope spreading".

In either case, it may be possible to restore tolerance in lupus patients by coupling antigenic oligonucleotides to a protein carrier such as gamma globulin (IgG). Calf-thymus DNA complexed to gamma globulin has been reported to reduce anti-DNA antibody formation.

Therapeutic Uses of Oligos Containing GCG Trinucleotide Sequences at or Near Both Termini

Based on their interaction with CREB/ATF, oligonucleotides containing GCG trinucleotide sequences at or near both termini have antiviral activity, independent of any antisense effect due to complementarity between the oligonucleotide and the viral sequence being targeted. Based on this activity, an effective amount of inhibitory oligonucleotides can be administered to a subject to treat or prevent a viral infection.

EXAMPLES

Example 1: Effects of ODNs on B Cell Total RNA Synthesis and Cell Cycle

B cells were purified from spleens obtained from 6-12 wk old specific
5 pathogen free DBA/2 or BXSB mice (bred in the University of Iowa animal care facility; no
substantial strain differences were noted) that were depleted of T cells with anti-Thy-1.2 and
complement and centrifugation over lympholyte M (Cedarlane Laboratories, Hornby,
Ontario, Canada) ("B cells"). B cells contained fewer than 1% CD4⁺ or CD8⁺ cells. 8x10⁴
10 B cells were dispensed in triplicate into 96 well microtiter plates in 100 µl RPMI containing
10% FBS (heat inactivated to 65°C for 30 min.), 50 µM 2-mercaptoethanol, 100 U/ml
penicillin, 100 µg/ml streptomycin, and 2 mM L-glutamate. 20 µM ODN were added at the
start of culture for 20 h at 37°C, cells pulsed with 1 µCi of ³H uridine, and harvested and
counted 4 hr later. Ig secreting B cells were enumerated using the ELISA spot assay after
15 culture of whole spleen cells with ODN at 20 µM for 48 hr. Data, reported in Table 1,
represent the stimulation index compared to cells cultured without ODN. Cells cultured
without ODN gave 687 cpm, while cells cultured with 20 µg/ml LPS (determined by titration
to be the optimal concentration) gave 99,699 cpm in this experiment. ³H thymidine
incorporation assays showed similar results, but with some nonspecific inhibition by
thymidine released from degraded ODN (Matson. S and A.M. Krieg (1992) Nonspecific
20 suppression of ³H-thymidine incorporation by control oligonucleotides. *Antisense Research
and Development* 2:325).

For cell cycle analysis, 2 X 10⁶ B cells were cultured for 48 hr. in 2 ml tissue
culture medium alone, or with 30 µg/ml LPS or with the indicated phosphorothioate modified
25 ODN at 1 µM. Cell cycle analysis was performed as described in (Darzynkiewicz, Z. et al.,
Proc. Natl. Acad. Sci. USA 78:2881 (1981)).

To test the mitogenic effects of CpG ODN on human cells, peripheral blood
monocyte cells (PBMCs) were obtained from two patients with chronic lymphocytic
30 leukemia (CLL), a disease in which the circulating cells are malignant B cells. Cells were
cultured for 48 hrs and pulsed for 4 hours with tritiated thymidine as described above.

Example 2: Effects of ODN on Production of IgM from B cells

Single cell suspensions from the spleens of freshly killed mice were treated
35 with anti-Thy1, anti-CD4, and anti-CD8 and complement by the method of Leibson et al., *J.
Exp. Med.* 154:1681 (1981)). Resting B cells (<0.2% T cell contamination) were isolated
from the 63 - 70% band of a discontinuous Percoll gradient by the procedure of DeFranco et
al, *J. Exp. Med.* 155:1523 (1982). These were cultured as described above in 30 µM ODN or

20 µg/ml LPS for 48 hr. The number of B cells actively secreting IgM was maximal at this time point, as determined by ELISpot assay (Klinman, D.M. et al. *J. Immunol.* 144:506 (1990)). In that assay, B cells were incubated for 6 hrs on anti-Ig coated microtiter plates. The Ig they produced (>99% IgM) was detected using phosphatase-labelled anti-Ig (Southern Biotechnology Associated, Birmingham, AL). The antibodies produced by individual B cells were visualized by addition of BCIP (Sigma Chemical Co., St. Louis MO) which forms an insoluble blue precipitate in the presence of phosphatase. The dilution of cells producing 20 - 40 spots/well was used to determine the total number of antibody-secreting B cells/sample. All assays were performed in triplicate. In some experiments, culture supernatants were assayed for IgM by ELISA, and showed similar increases in response to CpG-ODN.

table 1

Example 3: B cell Stimulation by Bacterial DNA

DBA/2 B cells were cultured with no DNA or 50 µg/ml of a) *Micrococcus lysodeikticus*; b) NZB/N mouse spleen; and c) NFS/N mouse spleen genomic DNAs for 48 hours, then pulsed with ³H thymidine for 4 hours prior to cell harvest. Duplicate DNA samples were digested with DNase I for 30 minutes at 37 C prior to addition to cell cultures. E coli DNA also induced an 8.8 fold increase in the number of IgM secreting B cells by 48 hours using the ELISA-spot assay.

20

DBA/2 B cells were cultured with either no additive, 50 µg/ml LPS or the ODN 1; 1a; 4; or 4a at 20 uM. Cells were cultured and harvested at 4, 8, 24 and 48 hours. BXSb cells were cultured as in Example 1 with 5, 10, 20, 40 or 80 µM of ODN 1; 1a; 4; or 4a or LPS. In this experiment, wells with no ODN had 3833 cpm. Each experiment was performed at least three times with similar results. Standard deviations of the triplicate wells were <5%.

Example 4: Effects of ODN on natural killer (NK) activity

10 x 10⁶ C57BL/6 spleen cells were cultured in two ml RPMI (supplemented as described for Example 1) with or without 40 µM CpG or non-CpG ODN for forty-eight hours. Cells were washed, and then used as effector cells in a short term ⁵¹Cr release assay with YAC-1 and 2C11, two NK sensitive target cell lines (Ballas, Z. K. et al. (1993) *J. Immunol.* 150:17). Effector cells were added at various concentrations to 10⁴ ⁵¹Cr-labeled target cells in V-bottom microtiter plates in 0.2 ml, and incubated in 5% CO₂ for 4 hr. at 37°C. Plates were then centrifuged, and an aliquot of the supernatant counted for radioactivity. Percent specific lysis was determined by calculating the ratio of the ⁵¹Cr released in the presence of effector cells minus the ⁵¹Cr released when the target cells are

cultured alone, over the total counts released after cell lysis in 2% acetic acid minus the ^{51}Cr cpm released when the cells are cultured alone.

Example 5: In vivo studies with CpG phosphorothioate ODN

5 Mice were weighed and injected IP with 0.25 ml of sterile PBS or the indicated phosphorothioate ODN dissolved in PBS. Twenty four hours later, spleen cells were harvested, washed, and stained for flow cytometry using phycoerythrin conjugated 6B2 to gate on B cells in conjunction with biotin conjugated anti Ly-6A/E or anti-Ia^d (Pharmingen, San Diego, CA) or anti-Bla-1 (Hardy, R.R. et al., *J. Exp. Med.* 159:1169 (1984). Two mice
10 were studied for each condition and analyzed individually.

Example 6 Titration of Phosphorothioate ODN for B Cell Stimulation

B cells were cultured with phosphorothioate ODN with the sequence of control ODN 1a or the CpG ODN 1d and 3Db and then either pulsed after 20 hr with ^3H
15 uridine or after 44 hr with ^3H thymidine before harvesting and determining cpm.

Example 7 Rescue of B Cells From Apoptosis

WEHI-231 cells (5×10^4 /well) were cultured for 1 hr. at 37 C in the presence
20 or absence of LPS or the control ODN 1a or the CpG ODN 1d and 3Db before addition of anti-IgM ($1\mu\text{ml}$). Cells were cultured for a further 20 hr. before a 4 hr. pulse with $2\mu\text{Ci}$ /well ^3H thymidine. In this experiment, cells with no ODN or anti-IgM gave 90.4×10^3 by addition of anti-IgM. The phosphodiester ODN shown in Table 1 gave similar protection, though with some nonspecific suppression due to ODN degradation. Each experiment was
25 repeated at least 3 times with similar results.

Example 8 In vivo induction of IL-6

DBA/2 female mice (2 mos. old) were injected IP with $500\mu\text{g}$ CpG or control phosphorothioate ODN. At various time points after injection, the mice were bled. Two
30 mice were studied for each time point. IL-6 was measured by Elisa, and IL-6 concentration was calculated by comparison to a standard curve generated using recombinant IL-6. The sensitivity of the assay was 10 pg/ml . Levels were undetectable after 8 hr.

Example 9 Binding of B cell CREB/ATF to a radiolabelled doublestranded CRE probe (CREB)

35 Whole cell extracts from CH12.LX B cells showed 2 retarded bands when analyzed by EMSA with the CRE probe (free probe is off the bottom of the figure). The

CREB/ATF protein(s) binding to the CRE were competed by the indicated amount of cold CRE, and by single-stranded CpG ODN, but not by non-CpG ODN.

Equivalents

5 Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents of the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

- 29 -

SEQUENCE LISTING

(1) GENERAL INFORMATION:

5

(i) APPLICANT: Arthur M. Krieg, M.D.

(ii) TITLE OF INVENTION: IMMUNOMODULATORY OLIGONUCLEOTIDES

10

(iii) NUMBER OF SEQUENCES: 27

(iv) CORRESPONDENCE ADDRESS:

15

(A) ADDRESSEE: LAHIVE & COCKFIELD

(B) STREET: 60 STATE STREET, SUITE 510

(C) CITY: BOSTON

(D) STATE: MASSACHUSETTS

(E) COUNTRY: USA

(F) ZIP: 02109-1875

20

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk

(B) COMPUTER: IBM PC compatible

(C) OPERATING SYSTEM: PC-DOS/MS-DOS

(D) SOFTWARE: ASCII text

25

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: US

(B) FILING DATE:

(C) CLASSIFICATION:

30

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: ARNOLD, BETH E.

(B) REGISTRATION NUMBER: 35,430

(C) REFERENCE/DOCKET NUMBER: UIZ-013CP

35

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (617)227-7400

(B) TELEFAX: (617)227-5941

40

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

45

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

50

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

55

GGGGTCAACG TTCAGGGGGG

20

(2) INFORMATION FOR SEQ ID NO:2:

- 30 -

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 15 base pairs
 (B) TYPE: nucleic acid
5 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

15 GCTAGACGTT AGCGT 15

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 15 base pairs
20 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

30 GCTAGATGTT AGCGT 15

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:
35 (A) LENGTH: 15 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

40

(ix) FEATURE:
45 (A) NAME/KEY: misc_feature
 (B) LOCATION: 7
 (D) OTHER INFORMATION: "N indicates 5 methyl cytosine"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

50 GCTAGANGTT AGCGT 15

(2) INFORMATION FOR SEQ ID NO:5:

55 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 15 base pairs
 (B) TYPE: nucleic acid

- 31 -

(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: DNA

(ix) FEATURE:
10 (A) NAME/KEY: misc_feature
(B) LOCATION: 13
(D) OTHER INFORMATION: "N indicates 5 methyl cytosine"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:
15 GCTAGACGTT AGNGT 15

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:
20 (A) LENGTH: 15 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:
30 GCATGACGTT GAGCT 15

(2) INFORMATION FOR SEQ ID NO:7:

35 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

40 (ii) MOLECULE TYPE: DNA

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:
ATGGAAGGTC CAGCGTTCTC 20

(2) INFORMATION FOR SEQ ID NO:8:
50 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
55 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- 32 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

5 ATCGACTCTC GAGCGTTCTC 20

(2) INFORMATION FOR SEQ ID NO:9:

10 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: DNA

20 (ix) FEATURE:

(A) NAME/KEY: misc_feature

(B) LOCATION: 3

(D) OTHER INFORMATION: "N indicates 5 methyl cytosine"

25 (ix) FEATURE:

(A) NAME/KEY: misc_feature

(B) LOCATION: 10

(D) OTHER INFORMATION: "N indicates 5 methyl cytosine"

30 (ix) FEATURE:

(A) NAME/KEY: misc_feature

(B) LOCATION: 14

(D) OTHER INFORMATION: "N indicates 5 methyl cytosine"

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

ATNGACTCTN GAGNGTTCTC 20

(2) INFORMATION FOR SEQ ID NO:10:

40 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

45 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

50 (ix) FEATURE:

(A) NAME/KEY: misc_feature

(B) LOCATION: 3

(D) OTHER INFORMATION: "N indicates 5 methyl cytosine"

55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

ATNGACTCTC GAGCGTTCTC

20

(2) INFORMATION FOR SEQ ID NO:11:

- 5 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE: DNA

15 (ix) FEATURE:

- (A) NAME/KEY: misc_feature
 (B) LOCATION: 18
 (D) OTHER INFORMATION: "N indicates 5 methyl cytosine"

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

ATCGACTCTC GAGCGTTNTC

20

(2) INFORMATION FOR SEQ ID NO:12:

25

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: DNA

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

ATGGAAGGTC CAACGTTCTC

20

40 (2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

45

(ii) MOLECULE TYPE: DNA

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GAGAACGCTG GACCTTCCAT

20

55

(2) INFORMATION FOR SEQ ID NO:14:

- 34 -

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA
- 10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:
GAGAACGCTC GACCTTCCAT 20
- 15 (2) INFORMATION FOR SEQ ID NO:15:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
20 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: DNA
- 25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:
GAGAACGCTC GACCTTCGAT 20
- 30 (2) INFORMATION FOR SEQ ID NO:16:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
35 (B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: DNA
- 40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:
GAGCAAGCTG GACCTTCCAT 20
- (2) INFORMATION FOR SEQ ID NO:17:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
50 (B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: DNA
- 55

- 35 -

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 6
- (D) OTHER INFORMATION: "N indicates 5 methyl cytosine"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

GAGAANGCTG GACCTTCCAT

20

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 14
- (D) OTHER INFORMATION: "N indicates 5 methyl cytosine"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

GAGAACGCTG GACNTTCCAT

20

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

GAGAACGATG GACCTTCCAT

20

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- 36 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

5 GAGAACGCTC CAGCACTGAT

20

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

- 10 (A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

20 TCCATGTCGG TCCTGATGCT

20

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

30 (ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

35 TCCATGCTGG TCCTGATGCT

20

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

- 40 (A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
45 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- 50 (A) NAME/KEY: misc_feature
(B) LOCATION: 8
(D) OTHER INFORMATION: "N indicates 5 methyl cytosine"

55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

- 37 -

TCCATGTNGG TCCTGATGCT

20

(2) INFORMATION FOR SEQ ID NO:24:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
10 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

15 (ix) FEATURE:

- (A) NAME/KEY: misc_feature
(B) LOCATION: 12
(D) OTHER INFORMATION: "N indicates 5 methyl cytosine"

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

TCCATGTCGG TNCTGATGCT

20

(2) INFORMATION FOR SEQ ID NO:25:

- 25 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
30 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

TCCATGACGT TCCTGATGCT

20

40 (2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
45 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

TCCATGTCGG TCCTGCTGAT

20

55

(2) INFORMATION FOR SEQ ID NO:27:

- 38 -

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

GGGGTCAAGT CTGAGGGGGG

20

- 39 -

CLAIMS

- 5 1. An oligonucleotide comprising from about 2 to about 100 nucleotides and containing at least one unmethylated CpG dinucleotide.
2. The oligonucleotide of claim 1 which contains a sequence represented by the following formula:
- 10
$$5' \quad X_1X_2CGX_3X_4 \quad 3'$$
- wherein C and G are unmethylated, X_1 , X_2 , X_3 and X_4 are nucleotides and a GCG trinucleotide sequence is not present at or near the 5' and 3' termini.
- 15 3. The oligonucleotide of claim 2 having a phosphate backbone modification.
4. The oligonucleotide of claim 3 wherein the phosphate backbone modification is a phosphorothioate backbone modification.
- 20 5. The oligonucleotide of claim 1 comprising the following nucleotide sequence:
- 25
$$5' \text{ GGGGTCAACGTTGAGGGGGG } 3' \text{ (SEQ ID NO:1)}$$
6. The oligonucleotide of claim 5 having a phosphate backbone modification.
- 30 7. The oligonucleotide of claim 6 wherein the phosphate backbone modification is a phosphorothioate modification.
8. An oligonucleotide delivery complex comprising the oligonucleotide of claim 1 and a targeting means.
- 35 9. An oligonucleotide delivery complex of claim 8, wherein the targeting means is selected from the group consisting of cholesterol, virosome, liposome, lipid, and a target cell specific binding agent.

10. A pharmaceutical composition comprising the oligonucleotide of claim 9 and a pharmaceutically acceptable carrier.
- 5 11. A pharmaceutical composition comprising the oligonucleotide of claim 2 and a pharmaceutically acceptable carrier.
12. A method for activating a subject's B cells comprising contacting the B cells with an effective amount of the oligonucleotide of claim 1.
- 10 13. A method for activating a subject's B cells comprising contacting the B cells with an effective amount of the oligonucleotide of claim 2.
- 15 14. A method for activating a subject's natural killer cells comprising contacting the natural killer cells with an effective amount of the oligonucleotide of claim 1.
- 20 15. A method for activating a subject's natural killer cells comprising contacting the natural killer cells with an effective amount of the oligonucleotide of claim 2.
- 25 16. A method for treating, preventing or ameliorating an immune system deficiency in a subject comprising administering to the subject an effective amount of a pharmaceutical composition of claim 10.
- 30 17. A method for treating, preventing or ameliorating an immune system deficiency in a subject comprising the steps of:
- a) contacting lymphocytes obtained from the subject with a composition of claim 1 ex vivo, thereby producing activated lymphocytes; and
 - b) readministering the activated lymphocytes obtained in step a) to the subject.
- 35 18. A method for vaccinating a subject comprising administering to the subject a composition of claim 10 in conjunction with administration of a vaccine.

- 41 -

- 5 19. A method for treating a disease associated with an immune system activation in a subject comprising administering to the subject an effective amount of a neutral oligonucleotide alone or in conjunction with a pharmaceutically acceptable carrier.
- 10 20. A method of claim 19 wherein the disease associated with immune system activation is systemic lupus erythematosus.
21. A method of claim 19 wherein the disease associated with immune system activation is sepsis.
- 15 22. An improved method for performing antisense therapy comprising methylating CpG containing oligonucleotides prior to administration to a subject.
- 20 23. An improved method for in vivo diagnoses using oligonucleotide probes comprising methylating CpG containing oligonucleotides prior to administration to a subject.
- 25 24. An oligonucleotide which is capable of interfering with the activity of viral or cellular transcription factors and containing a consensus immunoinhibitory CpG motif represented by the formula:
- 5'GCGX_nGCG3'**
- wherein X = a nucleotide and n = in the range of 0-50.
- 30 25. An oligonucleotide of claim 24, wherein X is a pyrimidine.
26. An oligonucleotide of claim 25, wherein X_n is a CpG dinucleotide.
- 35 27. A method for treating or preventing a viral infection in a subject comprising administering to the subject an immunoinhibitory oligonucleotide of claim 24.

INTERNATIONAL SEARCH REPORT

Int'l Application No
PCT/US 95/01570

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07H21/00 C12Q1/68 A61K31/70

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07H C12Q A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP,A,0 468 520 (MITSUI TOATSU CHEMICALS, INC.) 29 January 1992 see the whole document ---	1-4, 10, 11, 19
A	JOURNAL OF VIROLOGY, vol.64, no.2, February 1990 pages 757 - 766 D.OTT ET AL. 'Sequence Analysis of Amphotropic and 10A1 Murine Leukemia Viruses : Close Relationship to Mink Cell Focus-Inducing Viruses.' see the whole document --- -/-	1

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *A* document member of the same patent family

Date of the actual completion of the international search

4 July 1995

Date of mailing of the international search report

11. 07. 95

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Scott, J

INTERNATIONAL SEARCH REPORT

Inte nal Application No
PCT/US 95/01570

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		Relevant to claim No.
Category *	Citation of document, with indication, where appropriate, of the relevant passages	
A	JOURNAL OF VIROLOGY, vol.46, no.1, April 1983 pages 204 - 211 R.REPASKE ET AL. 'Nucleotide Sequence of the env-Specific Segment of NFS-Th-1 Xenotropic Murine Leukemia Virus.' see the whole document ---	1
A	JOURNAL OF VIROLOGY, vol.53, no.2, February 1985 L.WOLFF ET AL. 'Sequence Comparisons of the Anemia- and Polycythemia-Inducing Strains of Friend' see the whole document ---	1
A	JOURNAL OF VIROLOGY, vol.64, no.11, November 1990 pages 5491 - 5499 A.C.MASSEY ET AL. 'Origin of Pathogenic Determinants of Recombinant Murine Leukemia Viruses : Analysis of Bxv-1-Related Xenotropic Viruses from CWD Mice.' see the whole document -----	1

1

information on patent family members

PCT/US 95/01570

Form PCT/ISA/210 (patent family annex) (July 1992)